

14

Uses of Flax**Linseed Oil
(High Linolenic Acid)****Raw & Refined
Bolted & Blown
Grinding Oils
Heat Boiled
(polymerized oils)**

Adhesives, i.e. manufacture of hardboard and fibreboard
Protective Coatings: paints, house paint primers, varnishes, lacquers, stains, alkyd resins, enamels, epoxidized oils
Floor Covering: linoleum
Oilcloth, tarpaulin and other coated fabrics
Patent leather
Industrial Chemicals: fatty acids, soap, glycerin
Printing inks, grinding oils, newsprint, core oils, caulking compounds, waterproofing compounds, mastic cements, foundry binders, brake lining, hardboard, shoe polish, herbicide and pesticide carrier
Antispalling and curing treatments for concrete
Tempering oil, bonding oil, and highly conjugated oils for hardboards

Seed**Specialty Edible Products:**

- cakes, muffins
- bread products (whole seed or flour)
- replacement for sesame seed in baking products
- cooked and dry cereals

Seed & Oil**Health Food:**

- source of linolenic acid
- to make pills
- inclusion in pharmaceutical mediator mutations

Linseed Meal**High protein animal feed**

Seed, oil, meal: Dairy and beef cattle, swine and poultry feed and also to increase levels of linolenic acid in food products, i.e. eggs, meat and milk

Pet Foods: seed and meal

Fish feed in fish culture: Seed and oil - source of linolenic acid required by growing fish, i.e. salmon

Table 1. Fatty Acid Composition of Major Seed Oils

	Fatty Acid Content %			
	Saturates	Oleic	Linoleic	Linolenic
Linseed Flax	9	20	13	58
Canola	6	58	26	10
Safflower	9	20	70	<1
Sunflower	11	20	69	-
Corn	13	25	61	1
Olive	14	77	8	1
Soybean	15	24	54	7
Peanut	18	48	34	-
Cottonseed	27	19	54	-

Source: Agricultural Handbook No. 8-4. Human Nutr. Inform. Serv., U.S. Dept. Agric., Washington, DC. 1979.

Table 2. Fatty Acid Composition of Linseed Flax and High Linolenic Flax (M 5791)

Fatty acid, % in oil		Linseed Flax	High Linolenic Flax
Palmitic	C16:0	5.4	4.5
Stearic	C18:0	3.1	2.2
Oleic	C18:1	17.1	9.7
Linoleic	C18:2	14.7	10.8
Linolenic	C18:3	59.6	72.8
Iodine value		196	217

Table 3. Linolenic Acid Content of Canadian Flaxseed by Province - 1992-1999*

	Manitoba	Saskatchewan	Alberta	Mean
1992	59.1	58.7	55.7	58.8
1993	57.9	61.0	60.7	59.7
1994	58.6	60.5	56.9	59.6
1995	55.8	59.4	60.8	58.1
1996	57.8	59.3	59.8	58.7
1997	58.8	57.4	58.9	58.0
1998	57.2	56.6	56.9	56.8
1999	60.4	59.4	59.0	59.6
Mean	58.2	59.0	58.6	58.7

Source: Quality of Western Canadian flaxseed, Grain Research Laboratory, Canadian Grain Commission, Winnipeg, MB. Linolenic acid determined by gas chromatography of methyl esters of fatty acids according to International Organization for Standardization; Animal and vegetable fats and oils 1505508:1990E.

Table 4. Linolenic Acid Content of Four Widely Grown Canadian Flax Cultivars in Flax Cooperative Tests* conducted at Eight Locations in Manitoba and Saskatchewan, 1995-1999.

	Manitoba				Saskatchewan			
	Morden	Portage la Prairie	Roseba nk	Brandon	Indian Head	Melfort	Saskatoon n	Scott
1995								
AC McDuff	52.5	53.8	53.5	49.0	57.2	56.2	54.9	56.2
CDC Normandy	54.0	55.7	55.8	52.8	60.2	56.8	56.5	58.7
Flanders	55.0	56.5	56.3	52.0	60.2	58.9	57.5	58.9
NorLin	53.8	55.4	54.5	53.1	60.0	56.7	56.5	58.9
Mean	53.8	55.4	55.0	51.7	59.4	57.1	56.3	58.2
1996								
AC McDuff	53.6	55.2	55.5	52.6	55.4	53.8	54.4	56.0
CDC Normandy	55.8	54.6	57.6	54.9	58.2	56.7	56.4	58.1
Flanders	57.3	58.3	58.0	54.3	57.8	56.2	56.8	558.7
NorLin	55.2	54.2	56.8	54.7	57.8	55.4	56.0	57.7
Mean	55.5	55.6	57.0	54.1	57.3	55.5	55.9	57.6
1997								
AC McDuff	53.3	55.4	56.0	52.2	49.7	53.1	52.1	50.6
CDC Normandy	53.0	55.0	58.0	52.1	52.9	51.6	55.1	54.3
Flanders	55.9	58.9	58.1	53.8	53.9	54.8	55.1	53.7
NorLin	52.9	56.8	57.7	53.1	51.9	52.5	55.4	53.9
Mean	53.8	56.5	57.4	52.8	52.1	53.0	54.4	53.1
1998								
AC McDuff	51.5	51.9	51.4	46.9	50.8	47.9	49.5	45.5
CDC Normandy	53.8	53.2	53.2	48.9	53.7	51.4	52.3	48.2
Flanders	52.6	54.3	54.6	49.7	53.5	52.7	53.0	47.8
NorLin	52.7	53.5	54.0	49.8	53.4	50.0	51.3	48.0
Mean	52.6	53.2	53.3	48.8	52.8	50.7	51.5	47.4

Table 4 (Continued)

	Manitoba				Saskatchewan			
	Morden	Portage la Prairie	Roseba nk	Brandon	Indian Head	Melfort	Saskatoon	Scott
							n	Mean
1999								
AC McDuff	57.1	58.8	59.2	58.4	59.8	53.4	51.6	52.0
CDC Normandy	59.9	59.0	60.0	58.2	60.1	53.5	56.1	55.5
Flanders	60.2	61.1	62.4	49.0	61.4	54.6	54.7	55.6
NorLin	59.4	59.0	59.8	58.8	59.4	53.3	55.4	54.9
Mean	59.1	59.6	60.3	58.6	60.2	53.7	54.4	54.5
Overall Mean	55.0	56.0	56.6	53.2	56.4	54.0	54.5	54.2

* Cooperative tests conducted by Prairie Registration Recommending Committee on Grain. Linolenic acid content determined by gas-liquid chromatography of the fatty acid esters using the method described by Daun et al., J. Amer. Oil Chemists' Society, 60, 1983.

Table 5. Linolenic Acid Content* of High Linolenic Flax (M 5791) in Replicated Trials in Manitoba, 1998 & 1999, in Comparison with Three Commercial Cultivars

	Morden	Burdick	Portage la Prairie
<u>1998</u>			
High Linolenic Flax	70.1		
AC Emerson	57.0		
Flanders	53.1		
NorLin	55.2		
<u>1999</u>			
<u>Test 1</u>			
High Linolenic Flax	70.9	71.9	72.7
AC Emerson	61.7	60.6	65.1
Flanders	60.4	60.6	62.5
NorLin	59.7	57.8	59.3
<u>Test 2</u>			
High Linolenic Flax	71.9	70.3	
AC Emerson	58.6	59.2	
Flanders	58.2	60.3	
NorLin	56.9	59.3	
<u>Test 3</u>			
High Linolenic Flax	72.0	71.3	
AC Emerson	60.5	61.6	
Flanders	60.0	60.9	
NorLin	58.6	58.2	

* Linolenic acid content determined by gas-liquid chromatography of the fatty acid esters using the method described by Daun et al., J. Amer. Oil Chemists' Society, 60, 1983.

Table 6. Linolenic Acid Content of High Linolenic Flax (M 5791) in Field Trials, 1998 & 1999

Location	Year	% Linolenic Content
Fisher Branch, MB	1998	72.7
		Field 1
	1999	72.7
		Field 2
Erickson, MB	1998	72.2
		Field 1
	1999	72.9
		Field 2
Gadsby, AB	1998	72.9
		Field 1
	1999	72.6
		Field 2